


UNIVERSITY PHYSICS

Chapter 6 GAUSS'S LAW

PowerPoint Image Slideshow




Flux

FIGURE 6.1




This chapter introduces the concept of flux, which relates a physical quantity and the area through which it is flowing. Although we introduce this concept with the electric field, the concept may be used for many other quantities, such as fluid flow. (credit: modification of work by “Alessandro”/Flickr)

FIGURE 6.3




The flux of an electric field through the shaded area captures information about the “number” of electric field lines passing through the area. The numerical value of the electric flux depends on the magnitudes of the electric field and the area, as well as the relative orientation of the area with respect to the direction of the electric field.

FIGURE 6.4




- (a) A planar surface S_1 of area A_1 is perpendicular to the electric field E . N field lines cross surface S_1 .
- (b) A surface S_2 of area A_2 whose projection onto the xz -plane is S_1 . The same number of field lines cross each surface.

FIGURE 6.5




The direction of the area vector of an open surface needs to be chosen; it could be either of the two cases displayed here. The area vector of a part of a closed surface is defined to point from the inside of the closed space to the outside. This rule gives a unique direction.

FIGURE 6.6




- (a) Two potential normal vectors arise at every point on a surface.
- (b) The outward normal is used to calculate the flux through a closed surface.
- (c) Only S_3 has been given a consistent set of normal vectors that allows us to define the flux through the surface.

FIGURE 6.7




Electric flux through a cube, placed between two charged plates. Electric flux through the bottom face ($ABCD$) is negative, because \vec{E} is in the opposite direction to the normal to the surface. The electric flux through the top face ($FGHK$) is positive, because the electric field and the normal are in the same direction. The electric flux through the other faces is zero, since the electric field is perpendicular to the normal vectors of those faces. The net electric flux through the cube is the sum of fluxes through the six faces. Here, the net flux through the cube is equal to zero. The magnitude of the flux through rectangle $BCKF$ is equal to the magnitudes of the flux through both the top and bottom faces.

FIGURE 6.8




A surface is divided into patches to find the flux.

FIGURE 6.9




Calculating the flux of E_0 through a rectangular surface.

FIGURE 6.10



Calculating the flux of E_0 through a closed cubic surface.

FIGURE 6.11



The electric field produces a net electric flux through the surface S .

FIGURE 6.12

Since the electric field is not constant over the surface, an integration is necessary to determine the flux.






FIGURE 6.13




A closed spherical surface surrounding a point charge q .

FIGURE 6.14



Flux through spherical surfaces of radii R_1 and R_2 enclosing a charge q are equal, independent of the size of the surface, since all E-field lines that pierce one surface from the inside to outside direction also pierce the other surface in the same direction.

FIGURE 6.15



Understanding the flux in terms of field lines.

- (a) The electric flux through a closed surface due to a charge outside that surface is zero.
- (b) Charges are enclosed, but because the net charge included is zero, the net flux through the closed surface is also zero.
- (c) The shape and size of the surfaces that enclose a charge does not matter because all surfaces enclosing the same charge have the same flux.


Gauss's Law

FIGURE 6.2



Karl Friedrich Gauss (1777–1855) was a legendary mathematician of the nineteenth century. Although his major contributions were to the field of mathematics, he also did important work in physics and astronomy.

FIGURE 6.16




The electric flux through any closed surface surrounding a point charge q is given by Gauss's law.

(a) Enclosed charge is positive.


(b) Enclosed charge is negative.

FIGURE 6.17




The flux through the Gaussian surface shown, due to the charge distribution, is $\Phi = (q_1 - q_2 - q_5)/\epsilon_0$.

FIGURE 6.19




Various Gaussian surfaces and charges.

A cubical Gaussian surface with various charge distributions.




Spheres


FIGURE 6.21



(a) Spherically symmetric




(b) Not spherically symmetric



(c) Spherically symmetric


Illustrations of spherically symmetrical and nonsymmetrical systems. Different shadings indicate different charge densities. Charges on spherically shaped objects do not necessarily mean the charges are distributed with spherical symmetry. The spherical symmetry occurs only when the charge density does not depend on the direction. In (a), charges are distributed uniformly in a sphere. In (b), the upper half of the sphere has a different charge density from the lower half; therefore, (b) does not have spherical symmetry. In (c), the charges are in spherical shells of different charge densities, which means that charge density is only a function of the radial distance from the center; therefore, the system has spherical symmetry.

FIGURE 6.22




The electric field at any point of the spherical Gaussian surface for a spherically symmetrical charge distribution is parallel to the area element vector at that point, giving flux as the product of the magnitude of electric field and the value of the area. Note that the radius R of the charge distribution and the radius r of the Gaussian surface are different quantities.

FIGURE 6.23




A spherically symmetrical charge distribution and the Gaussian surface used for finding the field (a) inside and (b) outside the distribution.

FIGURE 6.24



Electric field of a uniformly charged, non-conducting sphere increases inside the sphere to a maximum at the surface and then decreases as $1/r^2$. Here, E_R . The electric field is due to a spherical charge distribution of uniform charge density and total charge Q as a function of distance from the center of the distribution.

FIGURE 6.25



Electric field vectors inside and outside a uniformly charged sphere.

Cylinders


FIGURE 6.27



(a) Cylindrically symmetric



(b) Not cylindrically symmetric




(c) Not cylindrically symmetric



(d) Cylindrically symmetric

To determine whether a given charge distribution has cylindrical symmetry, look at the cross-section of an “infinitely long” cylinder. If the charge density does not depend on the polar angle of the cross-section or along the axis, then you have cylindrical symmetry. (a) Charge density is constant in the cylinder; (b) upper half of the cylinder has a different charge density from the lower half; (c) left half of the cylinder has a different charge density from the right half; (d) charges are constant in different cylindrical rings, but the density does not depend on the polar angle. Cases (a) and (d) have cylindrical symmetry, whereas (b) and (c) do not.

FIGURE 6.28



The electric field in a cylindrically symmetrical situation depends only on the distance from the axis. The direction of the electric field is pointed away from the axis for positive charges and toward the axis for negative charges.

FIGURE 6.29

The Gaussian surface in the case of cylindrical symmetry. The electric field at a patch is either parallel or perpendicular to the normal to the patch of the Gaussian surface.




FIGURE 6.30

A Gaussian surface surrounding a cylindrical shell.






FIGURE 6.31

A Gaussian surface within a cylindrical shell.




Planes

FIGURE 6.32

The components of the electric field parallel to a plane of charges cancel out the two charges located symmetrically from the field point P . Therefore, the field at any point is pointed vertically from the plane of charges. For any point P and charge q_1 , we can always find a q_2 with this effect.


FIGURE 6.33



A thin charged sheet and the Gaussian box for finding the electric field at the field point P . The normal to each face of the box is from inside the box to outside. On two faces of the box, the electric fields are parallel to the area vectors, and on the other four faces, the electric fields are perpendicular to the area vectors.


Metals

FIGURE 6.34




Polarization of a metallic sphere by an external point charge $+q$. The near side of the metal has an opposite surface charge compared to the far side of the metal. The sphere is said to be polarized. When you remove the external charge, the polarization of the metal also disappears.

FIGURE 6.35




In the presence of an external charge q , the charges in a metal redistribute. The electric field at any point has three contributions, from $+q$ and the induced charges $-\sigma_A$ and $+\sigma_B$. Note that the surface charge distribution will not be uniform in this case.

FIGURE 6.36



Electric charges on a conductor migrate to the outside surface no matter where you put them initially.

FIGURE 6.39



An infinitesimally small cylindrical Gaussian surface surrounds point P , which is on the surface of the conductor. The field is perpendicular to the surface of the conductor outside the conductor and vanishes within it.

A side view of an infinite conducting plate and Gaussian cylinder with cross-sectional area A .

FIGURE 6.40






FIGURE 6.42




An isolated conducting sphere.

FIGURE 6.43




Electric field of a positively charged metal sphere. The electric field inside is zero, and the electric field outside is same as the electric field of a point charge at the center, although the charge on the metal sphere is at the surface.

FIGURE 6.44



- (a) A charge inside a cavity in a metal. The distribution of charges at the outer surface does not depend on how the charges are distributed at the inner surface, since the E -field inside the body of the metal is zero. That magnitude of the charge on the outer surface does depend on the magnitude of the charge inside, however.
- (b) A charge outside a conductor containing an inner cavity. The cavity remains free of charge. The polarization of charges on the conductor happens at the surface.


FIGURE 6.45




The charges induced by two equal and opposite charges in two separate cavities of a conductor. If the net charge on the cavity is nonzero, the external surface becomes charged to the amount of the net charge.

End

EXERCISE 18




EXERCISE 30




EXERCISE 31




(a)



(b)



(c)



(d)

EXERCISE 35






FIGURE 6.46




A charge inside a cavity of a metal. Charges at the outer surface do not depend on how the charges are distributed at the inner surface since E field inside the body of the metal is zero.

EXERCISE 61




EXERCISE 62




EXERCISE 69




EXERCISE 72




EXERCISE 75




EXERCISE 77




EXERCISE 78




EXERCISE 82




EXERCISE 83




EXERCISE 84




EXERCISE 85




EXERCISE 86




EXERCISE 87




EXERCISE 88




EXERCISE 89



EXERCISE 93



EXERCISE 95





This OpenStax ancillary resource is © Rice University under a CC-BY 4.0 International license; it may be reproduced or modified but must be attributed to OpenStax, Rice University and any changes must be noted.