





# humility

/hjuˈmɪlɪti/

*noun*

the quality of having a modest or low view of one's importance.

"he needs the humility to accept that their way may be better"

*synonyms:* **modesty**, humbleness, modestness, **meekness**, lack of pride, lack of vanity, **diffidence**, unassertiveness

"he needs the humility to accept that their way may be better"


I do not care about jets.

Paraphrased from Sevil Salur

---

*Christine Nattrass (UTK), Jet Tools May 2019*

I want to learn about the QGP.



First part paraphrased from Sevil Salur. The cartoon is my fault.

# What should we measure?

# What should we measure?

## ***Jupiter and the Monkey***

Jupiter promised a royal reward to the one whose offspring should be deemed the handsomest.

The monkey came with the rest, and presented a flat-nosed, hairless, ill-featured young monkey.

A general laugh saluted her on the presentation of her son.

She resolutely said; "He is at least in the eyes of me, his mother, the dearest, handsomest, and most beautiful of all."



[http://aesopsfables.org/F9\\_Jupiter-and-the-Monkey.html](http://aesopsfables.org/F9_Jupiter-and-the-Monkey.html)

Abbreviated

# Jet measurement madlibs!

[Adjective] [noun] [observable] in [collision system]

Groomed  
Unfolded  
Event-engineered

Top quark  
B-jet  
Z-boson  
D meson

Correlations  
Di-jet asymmetries  
 $V_1, V_2, V_3, V_4 \dots$

Proton-lead  
Ultra-central collisions

# Jet measurement madlibs!

[Adjective] [noun] [observable] in [collision system]

Groomed  
Unfolded  
Event-engineered

Top quark  
B-jet  
Z-boson  
D meson

Correlations  
Di-jet asymmetries  
 $V_1, V_2, V_3, V_4 \dots$

Proton-lead  
Ultra-central collisions

**Groomed top quark di-jet asymmetries in ultracentral collisions**



# Jet measurement madlibs!

[Adjective] [noun] [observable] in [collision system]

Groomed  
Unfolded  
Event-engineered

Top quark  
B-jet  
Z-boson  
D meson

Correlations  
Di-jet asymmetries  
 $V_1, V_2, V_3, V_4 \dots$

Proton-lead  
Ultra-central collisions

**Groomed top quark di-jet asymmetries in ultracentral collisions**

**Unfolded Z-boson correlations in proton-lead**

# Jet measurement madlibs!

[Adjective] [noun] [observable] in [collision system]

Groomed  
Unfolded  
Event-engineered

Top quark  
B-jet  
Z-boson  
D meson

Correlations  
Di-jet asymmetries  
 $v_1, v_2, v_3, v_4, \dots$

Proton-lead  
Ultra-central collisions

**Groomed top quark di-jet asymmetries** in ultracentral collisions

**Unfolded Z-boson correlations** in proton-lead

**Groomed top jet  $v_4$**  in ultra-central collisions

# Jet measurement madlibs!

[Adjective] [noun] [observable] in [collision system]

Groomed  
Unfolded  
Event-engineered

Top quark  
B-jet  
Z-boson  
D meson

Correlations  
Di-jet asymmetries  
 $v_1, v_2, v_3, v_4, \dots$

Proton-lead  
Ultra-central collisions

**Groomed top quark di-jet asymmetries** in ultracentral collisions

**Unfolded Z-boson correlations** in proton-lead


**Groomed top jet  $v_4$**  in ultra-central collisions

**Event-engineered groomed b-jet top quark di-jet asymmetries** in ultra-central collisions

# Error bars matter

- Simple measurements with small error bars may offer stronger constraints than better/more intuitive/more clever observables with large error bars
- Sometimes it's surprising which measurements provide greatest constraints
- Examples:
  - Single particle spectra with small error bars constrain uncertainty on properties of the medium well [Phys.Rev. C89 \(2014\) no.3, 034917](#)
  - Relatively low Q, large x DIS and Drell-Yan data provide comparable constraints on Higgs production to LHC jet measurements [PoS DIS2018 \(2018\) 024](#)
- Correlations between uncertainties are also important.  
***Experiments should publish them.***

# Do we really learn anything from this?\*




**HUGE error bars!**


\*I really do like this measurement. It's nothing personal. I ask the same thing about the measurements I'm working on.

# Available data

RHIC Jet measurements



LHC Jet Measurements




**RHIC:** High- $p_T$  hadron correlations 52%, reconstructed jets 3%

**LHC:** High- $p_T$  hadron correlations 23%, reconstructed jets 40%

Analysis about 1 year old, some may disagree with classifications, but the gist holds



# What should we measure?

# What should we measure?








# What should we measure?







# What should we measure?





ALICE-PUB-93497




# What should we measure?



# What should we measure?





ALI-PREL-101616




ALI-PUB-93497



# What you don't see matters



# What you don't see matters




# What should we measure?

## Some of my answers


- Everything!
  - Martin Spousta: But we can't measure everything! We should focus!
  - Yen-jie Lee: We should measure things which are sensitive to different properties of the medium and which are human understandable.
- “Simple” measurements with small error bars
- Beware complicated measurements and black boxes.
- Have an open mind – your (my!) pet observable may not be the best

# Background






# Discussion of analysis technique in a typical jet paper/talk



# Signal vs Background

**Background**

**Signal**



# Beware black boxes!



# Beware black boxes!




# ATLAS

Constituent biases don't matter that much up here

## Background subtraction method:

- Iterative procedure
  - **Calorimeter jets:** Reconstruct jets with  $R=0.2$ .  $v_2$  modulated  $\langle \text{Bkgd} \rangle$  estimated by energy in calorimeters excluding jets with at least one tower with  $E_{\text{tower}} > \langle E_{\text{tower}} \rangle$
  - Track jets:** Use tracks with  $p_T > 4$  GeV/c
  - Calorimeter jets from above with  $E > 25$  GeV and track jets with  $p_T > 10$  GeV/c used to estimate background again.
- Calorimeter tracks matching one track with  $p_T > 7$  GeV/c or containing a high energy cluster  $E > 7$  GeV are used for analysis down to  $E_{\text{jet}} = 40$  GeV



ALI-DER-92552

But they do matter down here!

→ **Background subtraction in use by experiments is not benign!**

Phys. Lett. B 719 (2013) 220-241

# Definition of Jets in a Large Background

- **Organizers:** M. Connors, G. Milhano, C. Nattrass, R. Reed, S. Salur
- **Spectra conveners:** R. Kunnawalkam Elayavalli, Y. Mehtar-Tani (R. Bertens)
- **Correlation conveners:** J. Noronha-Hostler, J. Huang
- **Substructure conveners:** Y. Lee, Y. Chien





**Extensively discussed** the interplay between experimental techniques and theoretical calculations with the aim of **reaching an agreement\*** on the way forward for extracting jet measurements from large background events such as those in heavy ion collisions and high luminosity p-p or electron-ion collisions.

**\*Consensus on some points**


62 Registered but due to various visa & travel complications: 45 + several BNL employees attended.

# Recommendations




***Include anything correlated  
in definition of jet***

***Provide enough details to make  
comparisons between data and models***



***Reconsider role of  
collinear safety***



***Discuss and put effort into the problem***

# The Lisbon Accord

An aerial photograph of a narrow street in Lisbon, Portugal, showing a dense cluster of buildings with characteristic red-tiled roofs. In the background, a hill with a castle or fortress is visible under a bright sky.

- [Lisbon Accord](#) proposed that heavy ion analyses adopt RIVET in July 2014
- Workshop position paper will second this recommendation



# Problems with RIVET

- ~~Not all heavy ion functionality exists~~ **RIVET 2.7**
- Very few heavy ion analyses exist → **Use undergraduates!**
- Only takes in HEPMC 2.0 input → **Write a converter**
- No way to deal with fluctuations
- No option to fit functions
- Cannot run in batch mode

## Few heavy ion analyses in RIVET



**RIVET isn't  
useful for  
heavy ion  
analyses**

# Undergraduates!\*

JETSCAPE  
REU

Left to right: Ricardo Santos (Berea), James Neuhaus, Jerrica Wilson, Mariah McCreary, Christine Nattrass, Austin Schmier (UTK)

\*And one beginning graduate student with no programming experience.  
We acknowledge substantial support from the US NSF and the JETSCAPE Collaboration

# Course-based undergraduate research experience

CBE—Life Sciences Education, Vol. 15, No. 2 | Articles

 Free Access

## Early Engagement in Course-Based Research Increases Graduation Rates and Completion of Science, Engineering, and Mathematics Degrees

Stacia E. Rodenbusch, Paul R. Hernandez, Sarah L. Simmons, and Erin L. Dolan

Jennifer Knight, Monitoring Editor:

Published Online: 13 Oct 2017 | <https://doi.org/10.1187/cbe.16-03-0117>

 Sections  View Article

 Tools  Share

### Abstract

National efforts to transform undergraduate biology education call for research experiences to be an integral component of learning for all students. Course-based undergraduate research experiences, or CUREs, have been championed for engaging students in research at a scale that is not possible through apprenticeships in faculty research laboratories. Yet there are few if any studies that examine the long-term effects of participating in CUREs on desired student outcomes, such as graduating from college and completing a science, technology, engineering, and mathematics (STEM) major. One CURE program, the Freshman Research Initiative (FRI), has engaged thousands of first-year undergraduates over the past decade. Using propensity score–matching to control for student-level differences, we tested the effect of participating in FRI on students' probability of graduating with a STEM degree, probability of graduating within 6 yr, and grade point average (GPA) at graduation. Students who completed all three semesters of FRI were significantly more likely than their non-FRI peers to earn a STEM degree and graduate within 6 yr. FRI had no significant effect on students' GPAs at graduation. The effects were similar for diverse students. These results provide the most robust and best-controlled evidence to date to support calls for early involvement of undergraduates in research.

## **Phys 494 – Course-based Undergraduate Research Experience in Relativistic Heavy Ion Physics**

### **Instructor:**

Dr. Christine Nattrass

Office: SERF 609

Phone: 974-6211

Email: [christine.nattrass@utk.edu](mailto:christine.nattrass@utk.edu)

Office hours: TBA

**Teaching assistant:** N/A

**Class time & Location:** TR 12:40-1:55 SERF 210


### **Course Description:**

This course will incorporate undergraduates into a research project in high energy nuclear physics in a course setting. Each student will be responsible for implementing a heavy ion analysis in the program RIVET so that it can be used by the JETSCAPE collaboration to make comparisons between Monte Carlo models and data. Each student's project will be incorporated into a public software repository so that it is available to the field and, if possible, it will be validated by the relevant experiment and incorporated into the official RIVET software.



<https://i.ytimg.com/vi/mZnv6LXD9qs/maxresdefault.jpg>

# Analyses (almost) implemented in UTK copy of RIVET




<https://github.com/cnattras/rivet-hi>

Need to be finalized! Hold me to it!


# Problem: The background fluctuates

## Background

## Signal




...even in Monte Carlo!





# How to address fluctuations?

- Tag final state hadrons as part of signal or background and only put these into jet finder
  - May work OK but conceptually suboptimal
- Compare to data which have not been corrected for fluctuations in the background
  - Technically feasible but jet measurements would not be physically meaningful, would be sensitive to our understanding of soft physics
- Unfold Monte Carlo results
  - Treating MC just like data
  - Unfolding is highly non-trivial



# Conclusions

- We need to be humble
- Focus on reduced uncertainties
- Publish correlations between uncertainties!
- Treat MC like data
  - Implement analyses in RIVET!
  - Warning: fluctuations in full MC
- Beware black boxes



# humility

/hjuˈmɪlɪti/

*noun*



the quality of having a modest or low view of one's importance.

"he needs the humility to accept that their way may be better"

*synonyms:* **modesty**, humbleness, modestness, **meekness**, lack of pride, lack of vanity, **diffidence**, unassertiveness

"he needs the humility to accept that their way may be better"

# Progress



**WARNING: Very small sample size, highly biased results, need to coordinate with Przemyslaw, Jochen, & Jan Fiete to ensure code can eventually be committed. Fluctuations in combinatorial background not subtracted.**

Undergraduates [Mariah McCreary \(UTK\)](#), [James Neuhaus \(UTK\)](#), [Jerrica Wilson \(UTK\)](#), [Ricardo Santos \(Berea\)](#); Graduate student [Austin Schmier \(UTK\)](#); Post doc [Redmer Bertens \(UTK\)](#)

[Mariah and Ricardo implementing ALICE analyses](#), [James and Redmer providing key support](#).

Funding from JETSCAPE, Berea, and the UTK physics department.

# Summary of existing heavy ion analyses

- STAR
  - 6 analyses but none of them instantiated by STAR, some validated but none heavy ion
  - I am an “author” on one and I have no recollection of this whatsoever so I’m not sure how official they are
  - 2 implemented in my class this semester, 2 in progress
- PHENIX
  - 4 implemented in my class this semester
- ALICE
  - 10 pp analyses
  - 3 heavy ion analyses in RIVET-HI (multiplicity, charged spectra, dihadron correlations)
  - 2 jet spectra analyses and one jet v2 analysis in UTK copy of RIVET-HI, still require management of fluctuations
- ATLAS - 0
  - 1 jet spectra analysis in UTK copy of RIVET-HI, one in jet spectra analysis and one jet v2 analysis in progress, still require management of fluctuations
- CMS - 0
  - 1 jet fragmentation function analysis in UTK copy of RIVET-HI, jet fragmentation function analysis and jet spectra analysis in progress, still require management of fluctuations
- LHCb - 0