

# PropiMax<sup>®</sup> EC

## Section 18 Quarantine Exemption

### For Control of Rusts (*Phakopsora pachyrhizi*) in Soybean

#### For Distribution and Use Only in the State of Tennessee

#### ATTENTION

- **This Quarantine Exemption expires March 1, 2007.**
- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of pesticide application.
- The product PropiMax<sup>®</sup> EC fungicide, EPA Reg. No. 62719-346, (containing 41.8% propiconazole) manufactured by Dow AgroSciences LLC may be used.
- All applicable conditions and restrictions on the federally registered product label for PropiMax EC as well as the Section 18 use directions must be observed.

#### Directions for Use

Apply PropiMax EC at the rate of 4 - 8 fl oz per acre (0.113 - 0.225 lb a.i./acre). PropiMax EC may be applied using ground or aerial broadcast equipment or through sprinkler irrigation (chemigation) equipment. Best results are obtained when PropiMax EC is applied at the first sign of disease. If the disease is established use the higher rate in the rate range. Repeat applications may be made at an interval of 10 to 14 days.

**Chemigation:** Apply PropiMax EC in 1/4 to 1/2 inches of sprinkler irrigation. Follow recommendations for application rates, timing and frequency of application in crop-specific directions for use (above).

#### Restrictions:

- Use of PropiMax EC under this Quarantine Exemption will not be allowed until the time that soybean rust has been determined to be present in the continental United States by the United States Department of Agriculture/Animal and Plant Health Inspection Service (USDA/APHIS). Once USDA/APHIS has confirmed that soybean rust is present, all existing EPA-approved section 18s for soybean rust will be automatically released for implementation in Tennessee as the Tennessee Department of Agriculture feels is appropriate.
- **Do not** make more than 2 applications or apply more than 12 ounces (0.338 lb active ingredient) per acre per year.
- **Preharvest Interval:** The final application of PropiMax EC must occur no later than the R5 stage of growth.
- **Re-entry:** A restricted entry interval of 24 hours must be observed.
- **Do not** use hay, forage, or fodder from the soybean crop as any component of animal feed or bedding.
- **Rotational Crop Restrictions:** To avoid possible illegal residues, do not plant any rotational crop intended for food, grazing, or any component of animal feed or bedding within 105 days of application of PropiMax EC to the preceding crop, unless the rotational crop appears on the label for PropiMax EC.

Any adverse effects resulting from the use of PropiMax EC under this emergency exemption must be immediately reported to the Tennessee Department of Agriculture.

®Trademark of Dow AgroSciences LLC

R-142-019

File symbol: 04-TN-05

EPA-accepted: 06/21/04

Initial printing